

search notes terms

=> s vitis and LOX

L1 1 VITIS AND LOX

=> d l1 1

L1 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:118656 CAPLUS

DN 138:165736

TI Protein and cDNA sequences of lipoxygenase LOX genes from
Vitis vinifera and use

IN Descenzo, Richard A.; Irelan, Nancy A.

PA USA

SO U.S. Pat. Appl. Publ., 36 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003033627	A1	20030213	US 2001-978522	20011016
PRAI	US 2000-241220P	P	20001016		

=> s vinifera and LOX

L2 1 VINIFERA AND LOX

=> s LOX and food(w)additive

L3 4 LOX AND FOOD(W) ADDITIVE

=> d l3 1-4 ibib ab

L3 ANSWER 1 OF 4 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC. on STN

ACCESSION NUMBER: 1999:213429 BIOSIS

DOCUMENT NUMBER: PREV199900213429

TITLE: Kinetics of thermal inactivation of pea seed lipoxygenases
and the effect of additives on their thermostability.

AUTHOR(S): Busto, M. D.; Apenten, R. K. Owusu [Reprint author];
Robinson, D. S. [Reprint author]; Wu, Z.; Casey, R.;
Hughes, R. K.

CORPORATE SOURCE: Procter Department of Food Science, University of Leeds,
Woodhouse Lane, Leeds, LS2 9JT, UK

SOURCE: Food Chemistry, (May, 1999) Vol. 65, No. 3, pp. 323-329.
print.

CODEN: FOCHDJ. ISSN: 0308-8146.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 26 May 1999

Last Updated on STN: 26 May 1999

AB Mature pea seeds contain two major lipoxygenases (LOX)
isoenzymes designated LOX-2 and LOX-3. The thermal
inactivation of crude pea LOX and the recombinant LOX
(rLOX) were studied. Heat-inactivation plots for crude extracts of pea
LOX were linear from which thermodynamic activation parameters,
DELTAH, DELTAS and DELTAG have been estimated. The enzymatic activity
was relatively stable with a respective half-life (t1/2) at 60 degreeC of
54.2 min for LOX from pea (Pisum sativum L. cv. Birte) or 18.4
min for a mutant line lacking LOX-2. At 50degreeC the
thermostability of LOX-3 present in crude extracts of the mutant
strain (t1/2 = 66.8 min) was 90% greater than purified recombinant
LOX-3 (rLOX-3; t1/2 = 34.6 min). However, rLOX-3 was more
heat-stable than rLOX2. Both rLOX-3 and pea mutant line lacking
LOX-2 possessed considerable thermostability at 60degreeC (t1/2 =
16.5 min and 18.4 min, respectively). Even at the higher temperatures of

the same receiver. Recovery of the internal standard, N-nitrosoazetidine, added at the 10 ppb level, was 86.5%. In addition, a few samples of nitrite-treated salmon (lox) were also tested for N-nitroamines. The results show that the method is applicable to samples containing nitrite-treated fish and fish-derived products.

L3 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN
ACCESSION NUMBER: 1998:637609 CAPLUS
DOCUMENT NUMBER: 130:49589
TITLE: Natural antioxidant activity in some microalgal species
AUTHOR(S): Perelman, Alexander; Matsukawa, Ritsuko; Schlosberg, Michal; Cohen, Bat-Sheva; Fostik-Magyar, Claude; Dubinsky, Zvy
CORPORATE SOURCE: Department of Life Sciences, Bar-Ilan University, Ramat Gan, 52900, Israel
SOURCE: Israel Journal of Plant Sciences (1998), 46(2), 169-176
CODEN: IJUPEU; ISSN: 0792-9978
PUBLISHER: Laser Pages Publishing
DOCUMENT TYPE: Journal
LANGUAGE: English

AB At the present time there is an increasing awareness of the potential harmful effects of various chem. **food additives** used to delay spoilage. This resulted in the search for natural substances which may prevent oxidn. of various lipids, a main cause of the onset of rancidity. Various algae were examd. for their potential as sources of antioxidants. As part of that search, the amt. of double bonds in their constituent compds. and their antioxidant (AO) and lipoxxygenase (**LOX**) activities were followed in some microalgal species. The effect of light intensity, salinity, and temp. on the AO activity of lipid exts. from the microalga Navicula sp. was studied. Maximum AO activity was obtained in Navicula grown under low light intensity. AO activity of various algal fractions was studied both as inhibition of **LOX** activity and of linoleate autoxidn. Most of the **LOX** inhibitive activity was found in the membranal lipid-sol. fractions, whereas the inhibition of autoxidn. was assocd. with the water-sol. components of the cell cytoplasm.

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> FIL STNGUIDE

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	24.60	24.81

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-0.65	-0.65

FILE 'STNGUIDE' ENTERED AT 18:25:00 ON 17 DEC 2003
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Dec 12, 2003 (20031212/UP).

=> s modify and flavor and characteristics and food
1 MODIFY
0 FLAVOR
9 CHARACTERISTICS
32 FOOD

6 FOODS
33 FOOD
 (FOOD OR FOODS)
L4 0 MODIFY AND FLAVOR AND CHARACTERISTICS AND FOOD

=> file agricola biosis embase caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.72	25.53

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-0.65

FILE 'AGRICOLA' ENTERED AT 18:32:24 ON 17 DEC 2003

FILE 'BIOSIS' ENTERED AT 18:32:24 ON 17 DEC 2003

COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 18:32:24 ON 17 DEC 2003

COPYRIGHT (C) 2003 Elsevier Inc. All rights reserved.

FILE 'CAPLUS' ENTERED AT 18:32:24 ON 17 DEC 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

=> s modify and flavor and food

L5 146 MODIFY AND FLAVOR AND FOOD

=> duplicate remove l5

DUPLICATE PREFERENCE IS 'AGRICOLA, BIOSIS, EMBASE, CAPLUS'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n

PROCESSING COMPLETED FOR L5

L6 124 DUPLICATE REMOVE L5 (22 DUPLICATES REMOVED)

=> s l6 and unpredictable

L7 0 L6 AND UNPREDICTABLE

=> s l6 and unpredictable?

L8 0 L6 AND UNPREDICTAB?

=> s l6 and LOX

L9 0 L6 AND LOX

=> s l6 and lipoxygenase

L10 2 L6 AND LIPOXYGENASE

=> d l1- 1-2 ibib ab

'L5-' IS NOT A VALID FORMAT FOR FILE 'BIOSIS'

The following are valid formats:

The default display format is BIB.

ABS	-----	AB
ALL	-----	AN, DN, TI, AU, CS, PI, SO, DT, FS, LA, OS, ED, AB, NCL, CC, IT (CT, ST), GT, ORGN, RN, CN, GEN
BIB	-----	AN, DN, TI, AU, CS, PI, SO, DT, FS, LA, OS, ED
CBIB	-----	AN, compressed bibliographic information
DALL	-----	ALL, delimited for post-processing
IABS	-----	ABS, with text label
IALL	-----	ALL, indented with text labels
IBIB	-----	BIB, indented with text labels

FULL ESTIMATED COST	22.82	48.35
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-0.65

FILE 'STNGUIDE' ENTERED AT 18:35:47 ON 17 DEC 2003
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
 LAST RELOADED: Dec 12, 2003 (20031212/UP).

=> file agricola biosis embase caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.18	48.53
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-0.65

FILE 'AGRICOLA' ENTERED AT 18:37:32 ON 17 DEC 2003

FILE 'BIOSIS' ENTERED AT 18:37:32 ON 17 DEC 2003
 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 18:37:32 ON 17 DEC 2003
 COPYRIGHT (C) 2003 Elsevier Inc. All rights reserved.

FILE 'CAPLUS' ENTERED AT 18:37:32 ON 17 DEC 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

=> s lipoxxygenase and wine and flavor
 L11 6 LIPOXYGENASE AND WINE AND FLAVOR

=> d l11 1-6 ti

L11 ANSWER 1 OF 6 AGRICOLA Compiled and distributed by the National
 Agricultural Library of the Department of Agriculture of the United States
 of America. It contains copyrighted materials. All rights reserved.
 (2003) on STN

TI Analysis of foods and beverages, headspace techniques

L11 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN

TI Protein and cDNA sequences of **lipoxxygenase** LOX genes from Vitis
 vinifera and use

L11 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN

TI Volatile compounds c6 in **wines** and other alcoholic beverages

~~L11 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN~~

~~TI Impact of fermentation technology on the phenolic and volatile composition
 of German red **wines**~~

L11 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN

TI Significance of **lipoxxygenase** in fruits and vegetables

L11 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN

TI Enzymes occurring in the formation of six-carbon aldehydes and alcohols in
 grapes

WEST**Freeform Search****Database:**

US Patents Full-Text Database
 US Pre-Grant Publication Full-Text Database
 JPO Abstracts Database
 EPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Term:
Display: **Documents in Display Format:** **Starting with Number**
Generate: ☐ Hit List ☒ Hit Count ☐ Side by Side ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

Cases

Search History
DATE: Wednesday, December 17, 2003 [Printable Copy](#) [Create Case](#)
Set Name
 side by side
Query
Hit Count
 result set

DB=USPT,PGPB,DWPI; PLUR=YES; OP=OR

<u>L10</u>	vitis.ab. and lipoxxygenase	2	<u>L10</u>
<u>L9</u>	vitis and lipoxxygenase	46	<u>L9</u>
<u>L8</u>	xygenaseL7	0	<u>L8</u>
<u>L7</u>	Vitis	1605	<u>L7</u>
<u>L6</u>	lipoxxygenase.clm. and vitis	1	<u>L6</u>
<u>L5</u>	LOX.clm. and vitis	5	<u>L5</u>
<u>L4</u>	LOX.clm. and vitis.ab.	1	<u>L4</u>
<u>L3</u>	LOX.ab. and vitis.clm.	0	<u>L3</u>
<u>L2</u>	LOX.clm. and vitis.clm.	1	<u>L2</u>
<u>L1</u>	LOX.clm and vitis.clm.	0	<u>L1</u>

END OF SEARCH HISTORY